



TITLE:

Remarks on test function methods for
blowup of solutions to semilinear evolution
equations in sectorial domain (Theoretical
Developments to Phenomenon Analyses
based on Nonlinear Evolution Equations)

AUTHOR(S):

Sobajima, Motohiro

CITATION:

Sobajima, Motohiro. Remarks on test function methods for blowup of solutions to semilinear evolution equations in sectorial domain (Theoretical Developments to Phenomenon Analyses based on Nonlinear Evolution Equations). 数理解析研究所講究録 2019, 2121: 63-73

ISSUE DATE:

2019-07

URL:

<http://hdl.handle.net/2433/252163>

RIGHT:

Remarks on test function methods for blowup of solutions to semilinear evolution equations in sectorial domain

東京理科大学理工学部数学科 側島 基宏 (Motohiro Sobajima)
Department of Mathematics, Faculty of Science and Technology,
Tokyo University of Science

1. Introduction

This paper is a joint work with Dr. Masahiro Ikeda (Keio University/RIKEN) and a part of joint works [6] and [5]. In this paper we consider the following initial-boundary value problem

$$(1.1) \quad \begin{cases} \tau \partial_t^2 u(x, t) - \Delta u(x, t) + \partial_t u(x, t) = |u(x, t)|^p, & (x, t) \in \Omega \times (0, T), \\ u(x, t) = 0 & (x, t) \in \partial\Omega \times (0, T), \\ u(x, 0) = \varepsilon f(x), \quad \tau \partial_t u(x, 0) = \tau \varepsilon g(x), & x \in \Omega, \end{cases}$$

where $\Omega \subset \mathbb{R}^N$ ($N \in \mathbb{N}$) is an unbounded domain, for instance,

$$\Omega_1 = \mathbb{R}^N \setminus \overline{B(0, 1)}, \quad \text{or} \quad \Omega(\Sigma) = \{\rho\omega \in \mathbb{R}^N; \rho > 0, \omega \in \Sigma\}$$

with $\Sigma \subset S^{N-1}$ having smooth boundary. The region $\Omega(\Sigma)$ is so-called sectorial domain. The parameters $p \in (1, \frac{N}{(N-2)_+})$ and $\varepsilon > 0$ describe the effect of nonlinearity and the smallness of initial data, respectively. Finally, the constant $\tau \in \{0, 1\}$ switches the parabolicity and hyperbolicity of the problem (1.1).

The interest of the present paper is the lifespan of blowup solutions to (1.1) for small initial data. Therefore we first fix the pair $(f, \tau g)$, then we discuss blowup of solutions to the problem (1.1) with sufficiently small $\varepsilon > 0$.

The study of global existence and blowup of solutions to (1.1) has a long history. In the case $\tau = 0$, the problem

$$(1.2) \quad \begin{cases} \partial_t u(x, t) - \Delta u(x, t) = u(x, t)^p, & (x, t) \in \mathbb{R}^N \times (0, T), \\ u(x, 0) = u_0(x) \geq 0, & x \in \mathbb{R}^N \end{cases}$$

is initially studied in [2] to understand the effects of dimension and nonlinearity. In [2], it is proved that

- (i) if $1 < p < 1 + \frac{2}{N}$, then (1.2) does not have non-trivial global-in-time solutions.
- (ii) if $p > 1 + \frac{2}{N}$, then (1.2) possesses nontrivial global-in-time solutions for small initial data.

The exponent $p_F = 1 + \frac{2}{N}$ is called Fujita exponent. The same consequence as (i) for the critical case $p = p_F$ is given in Hayakawa [3], Sugitani [25] and Kobayashi–Sirao–Tanaka [11]. The lifespan estimate of solutions to (1.2) is also studied in Lee–Ni [13] by using the heat kernel and the maximum principle as

$$(\text{Lifespan of } u \text{ with } u_0 = \varepsilon f) \sim \begin{cases} C\varepsilon^{-(\frac{1}{p-1} - \frac{N}{2})^{-1}} & \text{if } 1 < p < p_F, \\ \exp(C\varepsilon^{-(p-1)}) & \text{if } p = p_F. \end{cases}$$

In the case $\tau = 1$, the problem forms a Cauchy problem of the usual damped wave equation

$$(1.3) \quad \begin{cases} \partial_t^2 u(x, t) - \Delta u(x, t) + \partial_t u(x, t) = |u(x, t)|^p, & (x, t) \in \mathbb{R}^N \times (0, T), \\ u(x, 0) = \varepsilon f(x), \quad \partial_t u(x, 0) = \varepsilon g(x), & x \in \mathbb{R}^N. \end{cases}$$

The blowup phenomena and estimates of the lifespan of solutions to (1.3) has been studied from the work of Li–Zhou [16]. They proved the blowup of solutions with upper lifespan estimates for $1 < p \leq p_F$ when $N = 1, 2$. The three dimensional problem with sharp upper lifespan estimates is proved by Nishihara [22]. For general case, Todorova–Yordanov [26] showed blowup of small solutions for $1 < p < p_F$. Zhang [29] derived blowup of small solutions in the critical case $p = p_F$. For the lifespan estimates for the critical case $p = p_F$, recently, Lai–Zhou [12] succeeded in proving the sharp upper estimates by applying the consideration in [13]. The precise lifespan estimates for semilinear damped wave equation (1.3) is the same as that of semilinear heat equation (1.2).

Similar study of respective problems for halved spaces $\mathbb{R}^k \times \mathbb{R}^{N-k}$, for exterior domains and for sectorial domains has been separately done in the literature (see e.g., Meier [20, 21], Levine–Meier [14, 15], Ikehata [7, 8, 9, 10], Ogawa–Takeda [23], Pinsky [24] and Wakasugi [28]).

We point out that most of the blowup solutions in various equations (like (1.2) and (1.3)) can be treated in the framework of Mitidieri–Pokhozhaev [17] and also their lifespan estimates can be given by this argument with small modification (see Mitidieri–Pokhozhaev [19, 18]). However, in the critical case $p = p_F$ in (1.2) and (1.3), their argument does not give sharp lifespan estimates (see also, Ikeda–Ogawa [4]).

The purpose of the present paper is to propose a new test function method from the viewpoint of ordinary differential inequalities with respect to the parameter.

Here we introduce the definition of weak solutions to (1.1) which is used in the present paper.

Definition 1. For $(f, \tau g) \in H_0^1(\Omega) \times L^2(\Omega)$, the function $u : \Omega \times [0, T) \rightarrow \mathbb{R}$ is called the weak solution of (1.1) with initial data (f, g) in $(0, T)$ if u belongs the following class

$$S_T = \begin{cases} C([0, T); H_0^1(\Omega)) \cap C((0, T); L^{2p}(\Omega)) & \text{if } \tau = 0, \\ C^1([0, T); L^2(\Omega)) \cap C([0, T); H_0^1(\Omega)) \cap C([0, T); L^{2p}(\Omega)) & \text{if } \tau = 1 \end{cases}$$

and satisfies $u(0) = f$, $\tau \partial_t u(0) = \tau g$ and for every $\varphi \in C^1(\overline{\Omega} \times [0, T))$ with $\varphi = 0$ on $\partial\Omega \times [0, T)$,

$$\int_{\Omega} (\tau g + f) \varphi \, dx + \int_0^T \int_{\Omega} |u|^p \varphi \, dx \, dt = \int_0^T \int_{\Omega} \nabla u \cdot \nabla \varphi \, dx \, dt - \int_0^T \int_{\Omega} (\tau \partial_t u + u) \partial_t \varphi \, dx \, dt.$$

Remark 1.1. Existence of local-in-time weak solutions to (1.1) in the sense verifies by the standard argument of mild solutions

$$\begin{cases} U(t) = e^{t\Delta_D} f + \int_0^t e^{(t-s)\Delta_D} |u(s)|^p \, ds & \text{in } L^2(\Omega) & \text{if } \tau = 0, \\ u(t) = e^{-t\mathcal{A}} U_0 + \int_0^t e^{-(t-s)\mathcal{A}} \mathcal{N}(U(s)) \, ds & \text{in } H_0^1(\Omega) \times L^2(\Omega) & \text{if } \tau = 1, \end{cases}$$

where $\mathcal{A}(u, v) = (-v, -\Delta_D u + v)$ and $\mathcal{N}(u, v) = (0, |u|^p)$. The domains of Δ_D and \mathcal{A} are given by $D(\Delta_D) = H^2(\Omega) \cap H_0^1(\Omega)$ and $D(\mathcal{A}) = D(\Delta_D) \times H_0^1(\Omega)$. In this argument we require $p \leq \frac{N}{N-2}$ (see e.g., Cazenave–Haraux [1]).

The main result of this paper could be Proposition 2.1, which provides a sufficient condition on the shape of domain and the parameter p for the blowup phenomena of small solutions to (1.1). In Section 2, a positive harmonic function satisfying Dirichlet boundary condition plays a crucial role. The profile of this function can be regarded as the one of the shape of Ω . In Section 3, we give a result of lifespan estimates of blowup solutions in specified domains (Propositions 3.1, 3.2 and 3.3) as corollaries of Proposition 2.1.

2. Analysis of blowup via a test function method

Here we assume that there exists a positive harmonic function Φ satisfying Dirichlet boundary condition, that is, $\Phi \in C(\overline{\Omega}) \cap C^\infty(\Omega)$ satisfies

$$(2.1) \quad \begin{cases} \Delta \Phi(x) = 0 & x \in \Omega, \\ \Phi(x) > 0 & x \in \Omega, \\ \Phi(x) = 0 & x \in \partial\Omega. \end{cases}$$

Moreover, existence of a nonnegative auxiliary function $w \in C^2(\overline{\Omega})$ satisfying that there exists $k > 0$ such that

$$(2.2) \quad |\nabla w(x)|^2 \leq k w(x), \quad |\Delta w| \leq k.$$

is required. Note that unboundedness of Ω is required to ensure the existence of Φ .

Remark 2.1. Here we give examples of the choices of the pair (Φ, w) as follows:

$$(\Phi, w) = \begin{cases} (1, |x|^2) & \text{if } \Omega = \mathbb{R}^N \ (N \in \mathbb{N}), \\ (x_N, |x|^2) & \text{if } \Omega = \{x = (x_1, \dots, x_N) \in \mathbb{R}^N ; x_N > 0\}, \\ (1 - |x|^{2-N}, (|x| - 1)^2) & \text{if } \Omega = \{x \in \mathbb{R}^N ; |x| > 1\} \ (N \geq 3), \\ (\log |x|, (|x| - 1)^2) & \text{if } \Omega = \{x \in \mathbb{R}^2 ; |x| > 1\}. \end{cases}$$

The following assertion is the essential tool of the present work, which is derived via a test function method with a harmonic function Φ satisfying Dirichlet boundary condition.

Proposition 2.1. *Suppose that there exists a pair (Φ, w) such that Φ and w satisfy (2.1) and (2.2), respectively. Assume that $(f, \tau g) \in H_0^1(\Omega) \times L^2(\Omega)$ and $(\tau g + f)\Phi \in L^1(\Omega)$ with*

$$(2.3) \quad c_0 = \int_{\Omega} (\tau g + f)\Phi \, dx > 0.$$

If the function

$$(2.4) \quad h(T) = \int_1^T \left(\int_{\{x \in \Omega; w(x) < R\}} \left(1 + \frac{|\nabla w \cdot \nabla \Phi|}{\Phi} \right)^{p'} \Phi \, dx \right)^{1-p} dR$$

diverges as $T \rightarrow \infty$, then the solution of (1.1) with $\varepsilon \ll 1$ blows up until $T = h^{-1}(C\varepsilon^{-(p-1)})$, where C is a positive constant depending only on $N, p, f, \tau g$.

Remark 2.2. Proposition 2.1 asserts that blowup phenomena of solutions to (1.1) with small initial data are governed by the relation between the structure of the positive harmonic function Φ satisfying Dirichlet boundary condition and the exponent of non-linearity p .

Proof of Proposition 2.1. Here we fix two kinds of functions $\eta \in C^\infty([0, \infty))$ and $\eta^* \in L^\infty((0, \infty))$ as follows, which will be used in the cut-off functions:

$$\eta(s) \begin{cases} = 1 & \text{if } s \in [0, 1/2] \\ \text{is decreasing} & \text{if } s \in (1/2, 1) \\ = 0 & \text{if } s \in [1, \infty), \end{cases} \quad \eta^*(s) = \begin{cases} 0 & \text{if } s \in [0, 1/2), \\ \eta(s) & \text{if } s \in [1/2, \infty). \end{cases}$$

For $p > 1$, we define for $R > 0$,

$$\begin{aligned} \psi_R(x, t) &= [\eta(s_R(x, t))]^{2p'}, & (x, t) &\in \mathbb{R}^N \times [0, \infty), \\ \psi_R^*(x, t) &= [\eta^*(s_R(x, t))]^{2p'}, & (x, t) &\in \mathbb{R}^N \times [0, \infty) \end{aligned}$$

with

$$s_R(x, t) = \frac{w(x)^2 + t}{R}.$$

We also set $P(R) = \text{supp } \psi_R \cap (\Omega \times [0, R])$. The second function is useful in the sense of the equality $\frac{d}{ds}(\eta^q(s)) = q[\eta^*(s)]^{q-1}\eta'(s)$. This kind of test functions with $w(x) = |x|^\alpha$ is introduced in Mitidieri–Pokhozhaev [17].

Let u be a weak solution of (1.1) with initial data $(\varepsilon f, \varepsilon g)$ in $(0, T_\varepsilon)$. Without loss of generality, we assume that $T_\varepsilon > R_*$ with $R_* \geq 1$ satisfying for every $R \geq R_*$

$$I_R = \int_{\Omega} (\tau g + f)\Phi \psi_R(\cdot, 0) \, dx \geq \frac{c_0}{2} > 0.$$

This is possible by a consequence of the dominated convergence theorem. For $R \in (R_*, T_\varepsilon)$, take $\varphi = \Phi\psi_R$ as a test function which satisfies Dirichlet boundary condition in view of (2.1). Noting that by (2.2),

$$\begin{aligned} & |\tau\Phi\partial_t^2\psi_R - \Phi\partial_t\psi_R - \Delta(\Phi\psi_R)| \\ & \leq \frac{2p'\tau}{R^2} \left(|\eta''(s_R)|\eta(s_R) + (2p' - 1)(\eta'(s_R))^2 \right) \Phi[\psi_R^*]^{\frac{1}{p}} + \frac{2p'}{R} |\eta'(s_R)|\eta(s_R) \Phi[\psi_R^*]^{\frac{1}{p}} \\ & \quad + \frac{2p'}{R} \left(|\eta''(s_R)|\eta(s_R)|\Delta w| + (2p' - 1)(\eta'(s_R))^2 \frac{|\nabla w|^2}{R} \right) \Phi[\psi_R^*]^{\frac{1}{p}} + \frac{2p'}{R} |\nabla w \cdot \nabla \Phi| [\psi_R^*]^{\frac{1}{p}} \\ & \leq \frac{C'}{R} \left(1 + \frac{|\nabla w \cdot \nabla \Phi|}{\Phi} \right) \Phi[\psi_R^*]^{\frac{1}{p}}, \end{aligned}$$

where C is a positive constant depending only on $N, p, \|\eta\|_{W^{2,\infty}((0,\infty))}$ and k , we see from the definition of weak solutions and Hölder's inequality that

$$\begin{aligned} (2.5) \quad & \varepsilon I_R + \int_0^R \int_\Omega |u|^p \Phi \psi_R \, dx \, dt \\ & = \int_0^R \int_\Omega u \left(\tau\Phi\partial_t^2\psi_R - \Phi\partial_t\psi_R - \Delta(\Phi\psi_R) \right) \, dx \, dt \\ & \leq C \left(R^{-p'} \iint_{P(R)} \Theta^{p'} \Phi \, dx \, dt \right)^{\frac{1}{p'}} \left(\int_0^R \int_\Omega |u|^p \Phi \psi_R^* \, dx \, dt \right)^{\frac{1}{p}}, \end{aligned}$$

where

$$\Theta(x) = 1 + \frac{|\nabla w(x) \cdot \nabla \Phi(x)|}{\Phi(x)}.$$

Remark that the previous computation requires $\Delta\Phi = 0$. Now we define

$$Y(R) = \int_0^R \left(\int_0^\rho \int_\Omega |u|^p \Phi \psi_\rho^* \, dx \, dt \right) \frac{d\rho}{\rho}, \quad R_* \leq R < T_\varepsilon.$$

Then as in [6, Lemma 3.9], we have $Y \in C^1([R_*, T_\varepsilon])$ and

$$Y'(R) = \frac{1}{R} \int_0^R \int_\Omega |u|^p \Phi \psi_R^* \, dx \, dt, \quad Y(R) \leq \int_0^R \int_\Omega |u|^p \Phi \psi_R \, dx \, dt.$$

On the other hand, by the definition of $P(R)$, we have

$$\iint_{P(R)} \Theta^{p'} \Phi \, dx \, dt \leq \int_0^R \int_{\{x \in \Omega; w(x) < R\}} \Theta^{p'} \Phi \, dx \, dt = R \int_{\{x \in \Omega; w(x) < R\}} \Theta^{p'} \Phi \, dx.$$

Therefore (2.5) is reduced to the following ordinary differential inequality with respect to the parameter R :

$$(2.6) \quad \left(\frac{c_0 \varepsilon}{2} + Y(R) \right)^p \leq C^p \left(\int_{\{x \in \Omega; w(x) < R\}} \Theta^{p'} \Phi \, dx \right)^{p-1} Y'(R).$$

Solving the above equation, we deduce

$$\begin{aligned}
0 &\leq \left(\frac{c_0 \varepsilon}{2} + Y(R) \right)^{1-p} \\
&\leq \left(\frac{c_0 \varepsilon}{2} + Y(R_*) \right)^{1-p} - (p-1) C^{-p} \int_{R_*}^R \left(\int_{\{x \in \Omega; w(x) < \rho\}} \Theta^{p'} \Phi \, dx \right)^{1-p} d\rho \\
&\leq \left(\frac{c_0 \varepsilon}{2} \right)^{1-p} - (p-1) C^{-p} \int_{R_*}^R \left(\int_{\{x \in \Omega; w(x) < \rho\}} \Theta^{p'} \Phi \, dx \right)^{1-p} d\rho.
\end{aligned}$$

If (2.4) is satisfied, then we have $R \leq R_\varepsilon$ such that

$$\left(\frac{c_0 \varepsilon}{2} \right)^{1-p} = (p-1) C^{-p} \int_{R_*}^{R_\varepsilon} \left(\int_{\{x \in \Omega; w(x) < \rho\}} \Theta^{p'} \Phi \, dx \right)^{1-p} d\rho.$$

Since $R \in [R_*, T_\varepsilon)$ is arbitrary, we obtain $T_\varepsilon \leq R_\varepsilon$. □

3. The result of the problem in particular cases

In this section we give blowup results for the following cases:

- the whole space case $\Omega = \mathbb{R}^N$
- the case $\Omega = \Omega_1 = \{x \in \mathbb{R}^N; |x| > 1\}$ ($N \geq 2$)
- the case $\Omega = \Omega(\Sigma) = \{\rho\omega \in \mathbb{R}^N; \rho > 0, \omega \in \Sigma\}$ with $\Sigma \subset S^{N-1}$ ($N \geq 2$)

as application of Proposition 2.1.

3.1 The case $\Omega = \mathbb{R}^N$

The following assertion is well-known as mentioned in Introduction.

Proposition 3.1 (The case of $\Omega = \mathbb{R}^N$). *Let $(f, \tau g) \in H^1(\mathbb{R}^N) \times L^2(\mathbb{R}^N)$ with*

$$\tau g + f \in L^1(\mathbb{R}^N), \quad \int_{\mathbb{R}^N} (\tau g + f) \, dx > 0.$$

If $1 < p \leq p_F$, then the solution u of (1.1) blows up until

$$T = \begin{cases} C\varepsilon^{-(\frac{1}{p-1} - \frac{N}{2})^{-1}} & \text{if } 1 < p < 1 + \frac{2}{N}, \\ \exp(C\varepsilon^{-(p-1)}) & \text{if } p = 1 + \frac{2}{N}. \end{cases}$$

Proof. As in Remark 2.1, we choose

$$\Phi(x) \equiv 1, \quad w(x) = |x|^2$$

which satisfy (2.1) and (2.2), respectively. Then the function h in (2.4) is given by

$$\begin{aligned} h(T) &= \int_1^T \left(\int_{\{x \in \mathbb{R}^N; w(x) < R\}} \left(1 + \frac{|\nabla w \cdot \nabla \Phi|}{\Phi} \right)^{p'} \Phi dx \right)^{1-p} dR \\ &= |B(0, 1)|^{1-p} \int_1^T R^{-\frac{N}{2}(p-1)} dR \\ &= \begin{cases} \frac{|B(0, 1)|^{1-p}}{1-\frac{N}{2}(p-1)} (T^{1-\frac{N}{2}(p-1)} - 1) & \text{if } 1 < p < 1 + \frac{2}{N} \\ |B(0, 1)|^{1-p} \log T & \text{if } p = 1 + \frac{2}{N}, \end{cases} \end{aligned}$$

where $|B(0, 1)| = \int_{B(0, 1)} dx$. Therefore Proposition 2.1 implies the desired upper bound for the lifespan of solution u . \square

3.2 Exterior problem $\Omega = \Omega_1$ ($N \geq 2$)

Next assertion is a blowup result of small solutions to (1.1) in the exterior domain Ω_1 .

Proposition 3.2 (The case of $\Omega = \Omega_1$). *Let $(f, \tau g) \in H_0^1(\Omega_1) \times L^2(\Omega_1)$ with*

$$(\tau g + f) \log |x| \in L^1(\Omega_1), \quad \begin{cases} \int_{\Omega_1} (\tau g + f)(1 - |x|^{2-N}) dx > 0, & \text{if } N \geq 3, \\ \int_{\Omega_1} (\tau g + f) \log |x| dx > 0, & \text{if } N = 2. \end{cases}$$

If $1 < p \leq p_F$, then the solution u of (1.1) blows up until

$$T = \begin{cases} C\varepsilon^{-(\frac{1}{p-1} - \frac{N}{2})^{-1}} & \text{if } N \geq 3, 1 < p < 1 + \frac{2}{N}, \\ \exp(C\varepsilon^{-(p-1)}) & \text{if } N \geq 3, p = 1 + \frac{2}{N}, \\ C(\varepsilon^{-1} \log(\varepsilon^{-1}))^{\frac{p-1}{2-p}} & \text{if } N = 2, 1 < p < 2, \\ \exp(\exp(C\varepsilon^{-(p-1)})) & \text{if } N = 2, p = 2. \end{cases}$$

Remark 3.1. The cases $N \geq 3$ and $N = 2, 1 < p < 2$ are known (See Pinsky [24]). The main contribution of the paper [5] is the two-dimensional critical case $N = p = 2$.

Proof of Proposition 3.2. First we treat the case $N \geq 3$. We choose

$$\Phi(x) = 1 - |x|^{2-N}, \quad w(x) = (|x| - 1)^2.$$

Then we easily see that Φ satisfies $\Delta \Phi = 0$ and

$$|\nabla w|^2 = 4w, \quad |\Delta w| \leq 2N + 4, \quad |\nabla w \cdot \nabla \Phi| \leq 2(N - 2)\Phi.$$

By direct computation the function h in (2.4) can be estimated as follows:

$$h(T) \geq C \int_1^T \left(\int_{\{x \in \mathbb{R}^N; |x| < 2R^{\frac{1}{2}}\}} dx \right)^{1-p} dR.$$

By Proposition 2.1, we obtain the same conclusion as the case of $\Omega = \mathbb{R}^N$.

If $N = 2$, then we change the choice of Φ as

$$\Phi = \log |x|.$$

Then noting that

$$|\nabla w \cdot \nabla \Phi| = 2 \frac{|x| - 1}{|x|} \leq 2 \log |x| \leq 2\Phi,$$

we deduce that the function h in (2.4) is estimated as follows:

$$\begin{aligned} h(T) &\geq C \int_1^T (R \log R)^{1-p} dR \\ &\geq \begin{cases} CT^{2-p}(\log T)^{1-p} & \text{if } 1 < p < 2, \\ C \log \log T & \text{if } p = 2. \end{cases} \end{aligned}$$

Proposition 2.1 implies the desired upper bound for the lifespan of solution u . \square

3.3 Problems in $\Omega(\Sigma) = \{\rho\omega \in \mathbb{R}^N ; \rho > 0, \omega \in \Sigma\}$ ($N \geq 2$)

Finally, we treat the case of sectorial domain $\Omega = \Omega(\Sigma)$. By using Friedrichs extension, we can find local-in-time weak solutions to (1.1) for every $(f, \tau g) \in H_0^1(\Omega(\Sigma)) \times L^2(\Omega(\Sigma))$. The condition on essential selfadjointness of Laplacian Δ endowed with domain

$$D = \{u \in C_0^\infty(\mathbb{R}^N \setminus \{0\}) ; u = 0 \text{ on } \partial\Omega(\Sigma) (= \Omega(\partial\Sigma))\}$$

is written in [6].

To state the result for sectorial domain, First we state the assertion for the first eigenvalue and eigenfunction of the Laplace–Beltrami operator in Σ endowed with Dirichlet boundary condition (see [27, Chapter IX] for detail).

Lemma 1. *The Laplace–Beltrami operator $-\Delta_\Sigma$ in $L^2(\Sigma)$ endowed with domain $H^2(\Sigma) \cap H_0^1(\Sigma)$ is selfadjoint and all resolvent operator of $-\Delta_\Sigma$ are compact. The first eigenvalue λ_Σ is nonnegative and simple, and the corresponding eigenfunction φ_Σ is positive in Σ . Moreover, λ_Σ is positive if and only if $\Sigma \neq S^{N-1}$.*

Here we define γ as a smallest root of the quadratic equation $\gamma^2 + (N-2)\gamma - \lambda_\Sigma = 0$. Then the positive harmonic function on $\Omega(\Sigma)$ satisfying Dirichlet boundary condition is given as follows.

Lemma 2. *Set*

$$\Phi_\Sigma(x) = |x|^\gamma \varphi_\Sigma\left(\frac{x}{|x|}\right), \quad x \in \Omega(\Sigma).$$

Then Φ_Σ satisfies

$$\begin{cases} \Delta \Phi_\Sigma(x) = 0 & x \in \Omega(\Sigma), \\ \Phi_\Sigma(x) > 0 & x \in \Omega(\Sigma), \\ \Phi_\Sigma(x) = 0 & x \in \partial\Omega(\Sigma), \\ x \cdot \nabla \Phi_\Sigma(x) = \gamma \Phi_\Sigma(x) & x \in \Omega(\Sigma). \end{cases}$$

According to the blowup result with upper lifespan estimates for the case of $\Omega = \Omega(\Sigma)$ is the following. The critical exponent for (1.1) in $\Omega(\Sigma)$ seems to depend on γ which comes from the first eigenvalue of Laplace–Beltrami operator $-\Delta_\Sigma$.

Proposition 3.3 (The case of $\Omega = \Omega(\Sigma)$). *Let $(f, \tau g) \in H_0^1(\Omega(\Sigma)) \times L^2(\Omega(\Sigma))$ with*

$$(\tau g + f)\Phi_\Sigma \in L^1(\Omega(\Sigma)), \quad \int_{\Omega(\Sigma)} (\tau g + f)\Phi_\Sigma dx > 0.$$

If $1 < p \leq p_F$, then the solution u of (1.1) blows up until

$$T = \begin{cases} C\varepsilon^{-(\frac{1}{p-1} - \frac{N+\gamma}{2})^{-1}} & \text{if } 1 < p < 1 + \frac{2}{N+\gamma}, \\ \exp(C\varepsilon^{-(p-1)}) & \text{if } p = 1 + \frac{2}{N+\gamma}. \end{cases}$$

Proof. It is verified by choosing $\Phi = \Phi_\Sigma$ and $w(x) = |x|^2$. □

References

- [1] T. Cazenave, A. Haraux, “An introduction to semilinear evolution equations,” Translated from the 1990 French original by Yvan Martel and revised by the authors. Oxford Lecture Series in Mathematics and its Applications **13**. The Clarendon Press, Oxford University Press, New York, 1998.
- [2] H. Fujita, *On the blowing up of solutions of the Cauchy problem for $u_t = \Delta u + u^{1+\alpha}$* , J. Fac. Sci. Univ. Tokyo Sect. I **13** (1966), 109–124.
- [3] K. Hayakawa, *On nonexistence of global solutions of some semilinear parabolic differential equations*, Proc. Japan Acad. **49** (1973), 503–505.
- [4] M. Ikeda, T. Ogawa, *Lifespan of solutions to the damped wave equation with a critical nonlinearity*, J. Differential Equations **261** (2016), 1880–1903.
- [5] M. Ikeda, M. Sobajima, *Remark on upper bound for lifespan of solutions to semilinear evolution equations in a two-dimensional exterior domain*. J. Math. Anal. Appl. **470** (2019), 318–326.
- [6] M. Ikeda, M. Sobajima, *Sharp upper bound for lifespan of solutions to some critical semilinear parabolic, dispersive and hyperbolic equations via a test function method*, Nonlinear Anal. **182** (2019), 57–74.
- [7] R. Ikehata, *A remark on a critical exponent for the semilinear dissipative wave equation in the one dimensional half space*, Differential Integral Equations **16** (2003), 727–736.
- [8] R. Ikehata, *Critical exponent for semilinear damped wave equations in the N -dimensional half space*, J. Math. Anal. Appl. **288** (2003), 803–818.
- [9] R. Ikehata, *New decay estimates for linear damped wave equations and its application to nonlinear problem*, Math. Methods Appl. Sci. **27** (2004), 865–889.
- [10] R. Ikehata, *Two dimensional exterior mixed problem for semilinear damped wave equations*, J. Math. Anal. Appl. **301** (2005), 366–377.

- [11] K. Kobayashi, T. Sirao, H. Tanaka, *On the growing up problem for semilinear heat equations*, J. Math. Soc. Japan **29** (1977), 407–424.
- [12] N.-A. Lai, Y. Zhou, *The sharp lifespan estimate for semilinear damped wave equation with Fujita critical power in higher dimensions*, J. Math. Pures Appl. (9) **123** (2019), 229–243.
- [13] T.-Y. Lee, W.-M. Ni, *Global existence, large time behavior and life span of solutions of a semilinear parabolic Cauchy problem*, Trans. Amer. Math. Soc. **333** (1992), 365–378.
- [14] H.A. Levine, P. Meier, *A blowup result for the critical exponent in cones*, Israel J. Math. **67** (1989), 129–136.
- [15] H.A. Levine, P. Meier, *The values of the critical exponent for reaction-diffusion equation in cones*, Arch. Rational Mech. Anal. **109** (1990), 73–80.
- [16] T.T. Li, Y. Zhou, *Breakdown of solutions to $\square u + u_t = |u|^{1+\alpha}$* , Discrete Contin. Dynam. Systems **1** (1995), 503–520.
- [17] E. Mitidieri, S.I. Pokhozhaev, *A priori estimates and the absence of solutions of nonlinear partial differential equations and inequalities*, (Russian) Tr. Mat. Inst. Steklova **234** (2001), 1–384; translation in Proc. Steklov Inst. Math. **234** (2001), 1–362.
- [18] E. Mitidieri, S.I. Pokhozhaev, *Lifespan estimates for solutions of some evolution inequalities*, (Russian) Differ. Uravn. **45** (2009), 1441–1451; translation in Differ. Equ. **45** (2009), 1473–1484.
- [19] E. Mitidieri, S. Pohozaev, *Towards a unified approach to nonexistence of solutions for a class of differential inequalities*, Milan J. Math. **72** (2004), 129–162.
- [20] P. Meier, *Blow up of solutions of semilinear parabolic differential equations*, J. Appl. Math. Phys. **39** (1988), 135–149.
- [21] P. Meier, *On the critical exponent for reaction-diffusion equations*, Arch. Rational Mech. Anal. **109** (1990), 63–71.
- [22] K. Nishihara, *L^p - L^q estimates for the 3-D damped wave equation and their application to the semilinear problem*, Seminar Notes of Math. Sci., 6, Ibaraki Univ., 2003, 69–83.
- [23] T. Ogawa, H. Takeda, *Non-existence of weak solutions to nonlinear damped wave equations in exterior domains*, Nonlinear Anal. **70** (2009), 3696–3701.
- [24] R. Pinsky, *The Fujita exponent for semilinear heat equations with quadratically decaying potential or in an exterior domain*, J. Differential Equations **246** (2009), 2561–2576.
- [25] S. Sugitani, *On nonexistence of global solutions for some nonlinear integral equations*, Osaka J. Math. **12** (1975), 45–51.
- [26] G. Todorova, B. Yordanov, *Critical exponent for a nonlinear wave equation with damping*, J. Differential Equations **174** (2001), 464–489.
- [27] N. Ja. Vilenkin, “Fonctions spéciales et théorie de la représentation des groupes,” Dunod Paris, 1969; English translation, “Special functions and the theory of group representation,” Trans. Math. Monographs **22**, Amer. Math. Soc., Providence, R.I., 1968.

- [28] Y. Wakasugi, *A remark on the critical exponent for the semilinear damped wave equation on the half-space*, to appear in the proceedings of the 11th ISAAC congress 2017, in the session Recent Progress in Evolution Equations-igpde2, [arXiv:1708.06429](#).
- [29] Q.S. Zhang, *A blow-up result for a nonlinear wave equation with damping: the critical case*, C. R. Acad. Sci. Paris Sér. I Math. **333** (2001), 109–114.